

**Universitat de Lleida**

Document downloaded from:

<http://hdl.handle.net/10459.1/65102>

The final publication is available at:

<https://doi.org/10.1016/j.agry.2018.09.004>

Copyright

cc-by-nc-nd, (c) Elsevier, 2018



Està subjecte a una llicència de [Reconeixement-NoComercial-SenseObraDerivada 4.0 de Creative Commons](https://creativecommons.org/licenses/by-nc-nd/4.0/)

**Carbon footprint of cropping systems with grain legumes and cover crops:  
A case-study in SW France**

Daniel Plaza-Bonilla<sup>abc\*</sup>, Irene Nogué-Serra<sup>ac</sup>, Didier Raffailac<sup>c</sup>, Carlos  
Cantero-Martínez<sup>a</sup> and Éric Justes<sup>cd</sup>

<sup>a</sup>Department of Crop and Forest Sciences, Associated Unit EEAD-CSIC, Agrotecnio,  
University of Lleida, 25198 Lleida, Spain.

<sup>b</sup>Departamento de Suelo y Agua, Estación Experimental de Aula Dei, Consejo Superior de  
Investigaciones Científicas (EEAD-CSIC), POB 13034, 50080 Zaragoza, Spain.

<sup>c</sup>AGIR, Université de Toulouse, INPT, INP-PURPAN, INRA, 31320, Auzeville, France.

<sup>d</sup>Present address: CIRAD, UMR SYSTEM, Univ. Montpellier, CIHEAM-IAMM, CIRAD,  
INRA, Montpellier SupAgro, 34060 Montpellier, France (eric.justes@cirad.fr)

\*Corresponding author: daniel.plaza@pvcf.udl.cat (Daniel Plaza-Bonilla)

## Abstract

Agriculture contributes to a significant proportion of global emissions of greenhouse gases (GHG) but can also participate in climate change mitigation. The introduction of legumes in crop rotations reduces the dependence on N fertilizers and may mitigate the carbon (C) footprint of cropping systems. The aim of this study was to quantify the C footprint of six low-input arable cropping systems resulting from the combination of three levels of grain legumes introduction in a 3-yr rotation (GL0: no grain legumes, GL1: 1 grain legume, GL2: 2 grain legumes) and the use of cover crops (CC) or bare fallow (BF) between cash crops, covering two rotation cycles (6 years). The approach considered external emissions, on-site emissions and soil organic carbon (SOC) stock changes, and prioritized (i) field observations and (ii) simulation of non-measured variables with the STICS model, rather than default emission factors. As expected, fertilizers accounted for 80-90% of external emissions, being reduced by 50% and 102% with grain legumes introduction in GL1-BF and GL2-BF, compared to the cereal-based rotation (GL0-BF). Cover crops management increased machinery emissions by 24-35% compared to BF. Soil nitrous oxide (N<sub>2</sub>O) emissions were low, ranging between 205 and 333 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in GL1-BF and GL0-BF, respectively. Nitrate leaching represented the indirect emission of 11.6 to 27.2 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in the BF treatments and 8.2 to 10.7 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in the CC treatments. Indirect emissions due to ammonia volatilization ranged between 8.4 and 41.8 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>. The introduction of grain legumes strongly influenced SOC changes and, consequently, the C footprint. In the BF systems, grain legumes introduction in the rotations led to a significant increase in the C footprint, because of higher SOC losses. Contrarily, the use of cover crops mitigated SOC losses, and lowered the C footprint. These results indicated the need of CC when increasing the number of grain legumes in cereal-based rotations. Despite the multiple known benefits of introducing grain legumes in

44 cropping systems our research highlights the need to consider soil organic carbon changes in  
45 environmental assessments.

46 **Keywords:** external emissions; greenhouse gases; nitrous oxide; on-site emissions; STICS  
47 model.

## 1. Introduction

Agricultural production contributes to a significant proportion of global emissions of greenhouse gases (GHG), which contribute to global warming. The emissions related to fertilizer production and application to crops, machinery use, and various soil processes represent the main mechanisms underlying GHG emission to the atmosphere from arable crop production (Gan *et al.*, 2012). In this context, the quantification of the carbon footprint is an appropriate tool to estimate the impact of crop production on climate (Knudsen *et al.*, 2014). The C footprint is defined as “the quantity of GHG expressed in terms of carbon dioxide equivalents emitted to the atmosphere by an individual, organization, process, product or event from within a specified boundary” (Pandey *et al.*, 2011).

During the last decades, agricultural production in western countries has relied strongly on the application of nitrogen (N) fertilizers. The availability of synthetic N facilitated the specialization of arable cropping systems on the production of cereals, and made European agriculture highly dependent on synthetic fertilizer-N. However, the mismanagement of this fertilizer, e.g. use of excessive rates and/or application at periods of low crop needs, leads to negative environmental impacts. Among other, nitrate pollution of groundwaters, atmospheric pollution from ammonia volatilization, and contribution to global warming due to nitrous oxide (N<sub>2</sub>O) emissions constitute main environmental risks derived from inadequate N fertilizer management in agriculture (Bouwman *et al.*, 2013). The diversification of cropping systems with the introduction of legumes as cash crops, cover crops or intercrops represents a key strategy to reduce N fertilizer needs at the crop- and rotation scale (e.g. Bedoussac *et al.*, 2015; Plaza-Bonilla *et al.*, 2017b). Crop rotation with legumes also leads to other agronomic and environmental benefits such as a break-crop effect, which encompasses a range of factors that enhance the production of the subsequent crop due to the improvement of growing conditions (Watson *et al.*, 2017). However, to maximize their benefits, the introduction of legumes

requires the adaptation of the cropping system (Plaza-Bonilla *et al.*, 2017b; Reckling *et al.*, 2015).

Currently, the environmental assessments of crop production, such as the C footprint face different limitations. Knudsen *et al.*, (2014) stressed the importance of analyzing the full crop rotation when quantifying the C footprint of low-input cropping systems. The statement of Knudsen *et al.*, (2014) was based on the high reliance of low-input systems on nutrient recycling and green manuring, where the different cash and cover crops of a rotation are interlinked. Different authors also highlighted the need to include soil organic carbon (SOC) changes in C footprint assessments, given the impact of crop rotations and management practices on SOC (Gan *et al.*, 2012; Knudsen *et al.*, 2014; Pandey and Agrawal, 2014). In this line, the inclusion of cover crops in cropping systems has been reported as a feasible strategy to increase the amount of soil organic C (Poeplau and Don, 2015), highlighting the need to include SOC changes in environmental assessments (Prechsl *et al.*, 2017). SOC change is a dynamic, equilibrium oriented process, with constant in- and outflows that depend on management and environmental conditions. Inherent to this perception is that a given stock of SOC can only be depleted once; changes in management would always leave the soil carbon and N cycles strive to a new equilibrium. Essential is thereby the time required to arrive at a new equilibrium and how to set up an experimental approach that captures the new equilibrium.

Another aspect of controversy is the use of global default values for estimating soil GHG emissions, given the extreme variability of pedoclimatic conditions and cropping systems in which crop production takes place, which leads to significant uncertainties in the calculation of some processes such as the emission of N<sub>2</sub>O from soils. Regarding to this, Barton *et al.* (2014) pointed out the need to use site-specific field-based measurements to assess greenhouse gases emissions from cropping systems including grain legumes. Consequently, the collection of empirical data and the use of biogeochemical models can help to increase the accuracy of

98 current GHG emissions assessments (Lares-Orozco *et al.*, 2016). The aim of this study was to  
99 quantify the impact of introducing grain legumes and cover crops on the C footprint of low-  
100 input cropping systems in an area of SW France. We hypothesized that the diversification of  
101 cropping systems with legumes would decrease the C footprint thanks to the savings in  
102 synthetic N fertilizer, while the use of cover crops would decrease the C footprint thanks to  
103 their positive impact on SOC by increasing the photosynthetic activity versus time according  
104 to calendar year. Cover crop residues allow adding C to the soil. The approach considered  
105 external emissions and on-site emissions. The impact of the inclusion of soil organic carbon  
106 (SOC) stock changes on the C footprint calculation was also assessed, as crop rotations and  
107 management practices play a major role on the capacity of soils to store C due to their effects  
108 on soil organic C mineralization and crop residues C inputs.

## 2. Materials and methods

### 2.1. Experimental design

A field experiment was established in 2003 in the Institut National de la Recherche Agronomique (INRA) in Auzeville (SW France, 43° 31' 42'' N, 1° 28' 56'' E), representative of the pedoclimatic conditions of the Garonne valley. The aim of the experiment was the design and assessment of different low-input innovative cropping systems based on the introduction of grain legumes and cover crops. The cropping systems mainly differed in the amount of synthetic N fertilizer required. Over the last three decades, mean annual rainfall, air temperature and potential evapotranspiration were 685 mm, 13.7°C, and 905 mm, respectively. Soil characteristics (0-30 cm depth) were analyzed at the beginning of the experiment: soil texture was clay loam, mean ( $\pm 1$  standard deviation) pH (H<sub>2</sub>O, 1:2.5) was  $7.0 \pm 0.5$ , CEC was  $18.1 \pm 3.6$  cmol<sup>+</sup> kg<sup>-1</sup>, organic C was  $8.7 \pm 1.0$  g kg<sup>-1</sup> and organic N was  $1.1 \pm 0.1$  g kg<sup>-1</sup>.

Six different cropping systems were compared being the result of the combination of three levels of grain legumes introduction in a 3-yr rotation (GL0: no grain legumes, GL1: 1 grain legume, GL2: 2 grain legumes) and the use of cover crops (CC) or bare fallow (BF) between cash crops (Fig. 1). The GL0 treatment consisted in a sorghum (*Sorghum bicolor* L.) – sunflower (*Helianthus annuus* L.) – durum wheat (*Triticum turgidum* L.) rotation, the GL1 treatment consisted in a sunflower – winter pea (*Pisum sativum* L.) – durum wheat rotation and the GL2 treatment consisted in a soybean (*Glycine max* L.) – spring pea – durum wheat rotation. Durum wheat was a common crop in the three rotations to act as an indicator of the carryover effect of the different cropping systems. Different cover crops were used on each cropping system. In the GL0-CC system cover crops were vetch (*Vicia sativa* L.) and a vetch – oat (*Avena sativa* L.) mixture aimed to increase soil nitrogen availability for subsequent crops and reduce mineral nitrogen applications. In GL1-CC winter pea was accompanied by previous and succeeding mustard (*Sinapis alba* L.) cover crops aimed at reducing nitrate losses, while durum



wheat was followed by a vetch-oat mixture to increase soil nitrogen availability. Finally, in GL2-CC mustard was used after spring pea and durum wheat to reduce nitrate leaching given the higher presence of legumes in the rotation (Fig. 1). The analytical framework is based on the comparison of different cropping systems, each including a rotation with different crops and a management adapted to the specific crops sequence e.g. through individual fertilizer and tillage choices. This approach aims to consider the different adaptations that need to be carried out for an optimum functioning of cropping systems as done usually by farmers: e.g. adapting management practices such as N fertilization, tillage, irrigation, etc. to the sequence of crops and cover crops of the rotation and their impact on soil water, C and N dynamics. The first cropping season is not taken into account in the present study to avoid effects of the previous management on the results. Therefore, two cycles of a 3-year rotation (2005-2007 and 2008-2010) are analyzed. Within each 3-year rotation, each crop was grown every year to account for interannual climatic variability. The experiment was replicated in two contiguous blocks to include variability in soil texture.

## *2.2. Crop management*

Soil management was based on conventional intensive tillage using farmer's machinery as is traditional in the area. A pass of rotary harrow was implemented before sowing. In some years, a pass of moldboard plow was used to control weeds mechanically, to reduce the use of herbicides. A pass of cultipacker was used to ensure good contact between the soil and cover crops seeds. A disk plow was used to incorporate cover crops into the soil. As the experiment aimed at reducing chemical inputs, the destruction of cover crops was always done mechanically when the cover was not destroyed by frost, by the usual soil tillage or by a specific operation such as chopping crop residues before soil plowing, without using herbicide. One or two passes of a camera steering system guiding a cultivator were implemented to control weeds between sorghum and sunflower rows depending on the presence of weeds. Small grain crops

were seeded with a commercial seed driller, while a pneumatic row planter was used for sorghum, sunflower and soybean. Average sowing dates of cash crops can be found in Table 1. Different cultivars were used over the duration of the experiment to reduce the susceptibility to pathogens.

Nitrogen fertilization (only for non-legume cash crops) was adapted each year according to the balance-sheet method, considering: (i) requirements of the crop, (ii) residual soil N, and (iii) N mineralization estimated using a predicted mineral N balance (Plaza-Bonilla *et al.*, 2017b). Nitrogen fertilizer was applied with a pneumatic precision applicator splitting the rate in two or three applications (i.e. before sowing, beginning of stem elongation and booting) of ammonium nitrate for durum wheat and one single application of urea before sowing for sunflower and sorghum (Table 2). Depending on the cropping system 3-4 phosphorus (P) applications were carried out during the experimental period as triple superphosphate (45%  $P_2O_5$ ) before sowing according to soil analysis. Soil potassium levels were enough to satisfy crop needs according to soil analyses. Cash crop protection was performed according to the principles of integrated pest management. In general, one herbicide application was performed to all crops after sowing while, in some cases, other applications were carried out according to the sanitary state of crops. One or two fungicide applications were performed in durum wheat and pea. Sorghum and soybean were irrigated with a hose reel irrigation system during summer because of high demand for plant transpiration and low rainfall. The irrigation water rate was calculated to reach a target of ca. 80% of the maximum evapotranspiration of the crop. The same irrigation rate was applied to the CC and BF systems. Cash crop harvest (Table 1) was carried out with a commercial harvesting machine. A subsample was taken to determine grain moisture. The amount of grain C produced on each cropping system was calculated assuming a common grain C concentration of 420 g kg<sup>-1</sup>, whereas grain N was calculated as the product between dry matter grain yield and grain N

concentration, quantified by Dumas' method analysis. All crop residues were incorporated into the soil. During the development of the field experiment data regarding crop management practices carried out (i.e. date, type of practice), machinery and/or implements used, 115-hp tractor fuel consumption per ha, and type and rate/dose of inputs were registered.

### 2.3. Quantification of the carbon footprint

In order to compare the carbon footprint of the different cropping systems, the boundaries were established at field level. External emissions, on-site emissions and SOC stocks changes versus time were considered using the approaches described below. Based on the findings of Jambert *et al.* (1997) who worked in an arable area of SW France close to the present experiment, methane fluxes were considered negligible, being the maximum emission values reported by the authors of  $8.4 \text{ g C-CH}_4 \text{ ha}^{-1} \text{ d}^{-1}$ .

External emissions comprised the production and transport of seeds, pesticides, N and P fertilizers and energy needed to pump irrigation water. The calculation involved the product between the amount of each input and its emission factor (EF) (Table 3). In the case of pesticides containing more than one active ingredient (a.i.) the EF was calculated considering the (i) EF for each a.i., (ii) dose of each a.i., and (iii) the sum of a factor ( $0.4 \text{ kg Ceq kg a.i.}^{-1}$ ) to account for the emissions linked to the formulation process (Lal, 2004).

On-site emissions comprised machinery emissions because of fuel consumption, soil  $\text{N}_2\text{O}$  direct emissions,  $\text{N}_2\text{O}$  indirect emissions due to nitrate leaching to groundwater and deposition of ammonia volatilized and carbon dioxide ( $\text{CO}_2$ ) emissions because of urea hydrolysis. To quantify machinery emissions a fuel emission factor of  $0.9 \text{ kg Ceq kg}^{-1}$  (Lal, 2004) was multiplied by the amount of fuel consumed by each implement pass on an hectare basis (Table 4). Soil  $\text{N}_2\text{O}$  emissions, nitrate leaching and ammonia volatilization were simulated with the STICS model which was satisfactorily evaluated with a predictive quality

of various types of outputs of water, C and N dynamic balances on this site (Plaza-Bonilla *et al.*, 2015). We used model efficiency to measure the agreement between simulated and observed values. This statistical criterion ranges between 1, indicating perfect match between simulated and observed values, and infinite negative value, when observed mean is a better predictor than the model. In this line, model efficiency was 0.64 and 0.37 indicating correct performance for simulating soil water and mineral N contents over the rooting depth (0-120 cm). Model efficiency was also satisfactory for other variables. It ranged between 0.73 and 0.80 for grain N concentration, crop N uptake and legumes N<sub>2</sub> fixation, and was 0.90 and 0.63 when simulating crops aerial biomass and grain yield, respectively (Plaza-Bonilla *et al.*, 2015). Moreover, statistical criteria, such as  $r^2$ , model efficiency, relative root mean square error and mean difference, corresponding to the comparison between cumulative observed and simulated N<sub>2</sub>O emissions of a subsequent version of this experiment were also found correct, with values of 0.40, 0.24, 45.6% and 0.1 kg N<sub>2</sub>O–N ha<sup>-1</sup>, respectively (Plaza-Bonilla *et al.*, 2017a).

The soil and crop model STICS (Brisson *et al.*, 1998, 2002, 2008) is a one-dimension daily-step model. It uses pedoclimatic and crop characteristics, and management practices as inputs for the simulation of growing seasons resulting on different crop and soil outputs. Climatic inputs were obtained on the site (<1% of 100 m far) at a daily scale. In the model, soil was divided in five different layers (i.e. 0-15, 15-30, 30-60, 60-90, 90-120 cm) of given key characteristics. Different parameters such as crop species, phenology, leaf area index (LAI) and sensibility to stresses like drought are the crop parameters needed by the model. Finally, some information related to crop management practices (e.g. date, amount of input, soil depth) is required as a model input. Management practices information was available in the database of the experiment. Soil N<sub>2</sub>O emissions as a result of nitrification and denitrification processes are simulated separately in the model and coupled by nitrate production by nitrification. In the model nitrification is proportional to soil ammonium content. This process is modulated by

water filled pore space (WFPS), soil temperature and pH. However, soil pH values in the experiment, ranging between 6.6 and 8.2, do not constraint nitrification. Denitrification is proportional to soil nitrate concentration and soil denitrification potential rate, which is modulated by soil pH and WFPS. The rate was checked from field samples and was found equivalent to 2 kg N<sub>2</sub>O-N ha<sup>-1</sup> d<sup>-1</sup> (Plaza-Bonilla *et al.*, 2017a). Ammonia volatilization is proportional to N fertilizer rate and depends on crop N uptake rate and soil pH. The model was parameterized and evaluated for crop growth and water and mineral N dynamics to estimate nitrate leaching (Plaza-Bonilla *et al.*, 2015) and for direct soil N<sub>2</sub>O emissions and ancillary variables (Plaza-Bonilla *et al.*, 2017a) under the conditions of the field experiment. Simulated nitrate leaching and ammonia volatilization were transformed to indirect N<sub>2</sub>O emissions taking into account the EF reported by the IPCC, i.e. 0.75% and 1%, respectively (IPCC, 2006). Afterwards, N<sub>2</sub>O was transformed to C<sub>eq</sub> taking into account a global warming potential of 265 (IPCC, 2013). Finally, CO<sub>2</sub> emissions resulting from urea hydrolysis were calculated using an EF of 0.2 kg CO<sub>2</sub>-C kg<sup>-1</sup> urea (IPCC, 2006).

The change in the stock of soil organic carbon (SOC) at the plow layer (0-30 cm depth) was quantified by analyzing SOC concentration and soil bulk density at the beginning of the experiment (2003) and after one (2006) and two (2009) complete rotation cycles. Organic C concentration was determined with a Leco-2000 analyzer (LECO, St. Joseph, MI, US).

SOC stocks annual change was quantified as the slope of the linear relationship between SOC and the year of sampling using a mixed linear model with random effect to account for differences between replicates on the intercept while the slope was considered to be equal for the different replications (i.e. fixed effect) (Plaza-Bonilla *et al.*, 2016). This analysis helped to consider possible differences in initial soil organic C values between plots. That value was considered to be the representative of the annual change for the duration considered here.

257           The different items were calculated as six year average of two successive three-year  
258   rotations for each cropping system. When not explicitly described in the text, data are reported  
259   as kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>. Finally, for each cropping system, the carbon footprint was divided by  
260   the amount of grain C and grain N produced to obtain an indicator relating the environmental  
261   and agronomic dimensions. Analyses of variance were performed using the JMP 13 Pro  
262   statistical package (SAS Institute Inc, 2017) for a completely randomized design with blocks  
263   with cropping system and year and their interaction as fixed effects and block as random effect.

### 3. Results

#### 3.1. Environmental conditions and cropping systems productivity.

Precipitation differed greatly among the six cropping seasons studied, with values below the 30-yr average (685 mm) in the first four seasons (Fig. 2). Differently, the 2008-2009 season was characterized by a wetter autumn than the average and high precipitation values in April 2009. Mean annual temperature was close to the 30-yr average, 13.7°C, with values ranging between 13.3 and 14.2 °C.

Durum wheat grain yield presented a quite narrow range from 4974 to 5458 kg ha<sup>-1</sup>, without significant differences between cropping systems in the period studied (2005-2010) ( $P = 0.07$ ) (Table 5) due to a large soil water content at field capacity which allowed buffering precipitation water deficit at late development stages. Similarly, the incorporation of cover crops in the rotations did not affect significantly the grain yield of the other cash crops studied (with  $P$  values of 0.33, 0.51, 0.71, 0.66 and 0.75 for sorghum, sunflower, winter and spring pea and soybean, respectively) (Table 5).

#### 3.2. External emissions.

Greater annual external emissions were quantified for the cropping systems without grain legumes compared to GL1 and GL2, reaching values of 1362 and 1358 kg CO<sub>2</sub>eq ha<sup>-1</sup> yr<sup>-1</sup> in GL0-BF and GL0-CC, respectively (Fig. 3). External emissions related to fertilizers represented between 47 and 62% of the entire C footprint excluding SOC changes (Table S1). The adaptation of crop N fertilization in each cropping system had a great impact on external emissions. GL1-BF and GL2-BF lowered by 50% and 102% the external emissions related to N fertilizers, respectively, compared to GL0-BF. The reduction was smaller in the systems with cover crops, attaining a reduction of 40% and 70% in GL1-CC and GL2-CC compared to GL0-CC, respectively. Emissions related to N fertilizers accounted for 90% of external emissions in

GL0-BF, GL0-CC, GL1-BF, and GL1-CC, while that value was reduced by 80% in GL2-BF and GL2-CC. Differently, the proportion of emissions due to energy consumption for irrigation was higher for GL2-BF and GL2-CC systems reaching 12% of the external emissions, while ranged between 0 and 6% for the rest of treatments. The emissions related to seeds and pesticides only represented 5% and 1% of the external emissions, respectively, as an average of the different cropping systems compared. Durum wheat seeds represented the greatest emissions when compared with the rest of cash crops, due to the amount of input used (i.e. sowing rate) and N fertilizer requirements to produce this crop. In the cropping systems with cover crops, the use of vetch and vetch-oat mixtures particularly increased the indirect emissions related to seeds given their higher sowing rate compared to mustard. However, energy consumption for irrigation, and indirect emissions related to seeds and pesticides had a low contribution on the C footprint excluding SOC, representing a 4, 3 and 1%, respectively, as an average of the six cropping systems compared (Table S1).

### 3.3. On-site emissions.

#### 3.3.1. Machinery emissions and volatilized CO<sub>2</sub> from urea hydrolysis.

As an average of crop sequences and years, machinery emissions were similar between crop rotations without cover crops being the values 232, 250 and 245 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in the GL0-BF, GL1-BF and GL2-BF cropping systems, respectively (Fig. 4). The increase in the use of machinery for the establishment, chopping and incorporation by soil tillage of cover crops led to greater machinery emissions in GL0-CC, GL1-CC and GL2-CC, with values of 288, 337 and 317 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup>, respectively (Fig. 4). In average of cropping systems, machinery use represented between 11 and 23% of the C footprint excluding SOC changes (Table S1).



Carbon dioxide emission as a byproduct of urea hydrolysis was only accounted for sunflower and sorghum fertilization in GL0-BF and GL0-CC cropping systems and in 2009 for sunflower in GL2-BF and GL2-CC (Table 2). The resulting mean emissions of this process were 67 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> for the cropping systems without legumes and 3.0 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> for the cropping systems with one grain legume (Fig. 4), being a 3% of the C footprint excluding SOC changes (Table S1).

### 3.3.2. Simulation of soil N<sub>2</sub>O direct emissions and N<sub>2</sub>O indirect emissions.

As an average of the three crop sequences, simulated cumulative soil N<sub>2</sub>O emissions during the six years (from 2005 to 2010) were 4.8, 2.9 and 3.8 kg N<sub>2</sub>O-N ha<sup>-1</sup> for the GL0-BF, GL1-BF and GL2-BF. In the case of the counterparts with cover crops, the values were 3.7, 4.2 and 3.9 kg N<sub>2</sub>O-N ha<sup>-1</sup> for GL0-CC, GL1-CC and GL2-CC, respectively. In average of the six cropping systems, direct soil N<sub>2</sub>O emissions would represent between 13 and 20% of the C footprint excluding SOC changes. According to the STICS model simulations, soil N<sub>2</sub>O emissions were highly dynamic and dependent of the combination between the crop and the cropping seasons (Fig. 5). Peaks of greatest magnitude were simulated mainly after N application to summer non-legume crops (mainly sorghum and, with less importance, sunflower) in the most N-dependent cropping systems (i.e. GL0-BF and GL0-CC) (Fig. 5a-c). Moreover, in some cases, the incorporation of low C:N ratio biomass from cover crops increased simulated soil N<sub>2</sub>O emissions, such the case of vetch-oat mixture in GL1-CC (Fig. 5d). Except for durum wheat, the rest of crops present in GL2-BF and GL2-CC, the least N-dependent systems, led to small magnitude soil N<sub>2</sub>O peaks, although the basal emission of this gas from the soil did not decreased the cumulative values (Fig. 5g-i).

Simulated nitrate leaching in the different cropping systems compared was reported in a previous publication (Plaza-Bonilla *et al.*, 2015). During the six years analyzed the amount

of N lost as leaching was estimated to be 22.3, 32.0 and 52.2 kg N ha<sup>-1</sup> in GL0-BF, GL1-BF and GL2-BF, respectively, while the values were reduced to 15.7, 20.5 and 18.0 kg N ha<sup>-1</sup> in GL0-CC, GL1-CC and GL2-CC, respectively, as an average of the three possible crop sequences. These values represented the indirect emission of 11.6, 16.7 and 27.2 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in the GL0-BF, GL1-BF and GL2-BF treatments, respectively, and 8.2, 10.7 and 9.4 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in GL0-CC, GL1-CC and GL2-CC, respectively. According to the simulations performed, during the six years analyzed ammonia volatilization from fertilizer-N amounted to 52.7, 12.1 and 12.6 kg NH<sub>3</sub>-N ha<sup>-1</sup> in GL0-BF, GL1-BF and GL2-BF treatments, respectively, and to 60.3, 15.6 and 15.3 kg NH<sub>3</sub>-N ha<sup>-1</sup> in GL0-CC, GL1-CC and GL2-CC treatments, respectively. These values represented the indirect emission of 36.6, 8.4 and 8.8 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in the GL0-BF, GL1-BF and GL2-BF treatments, respectively, and 41.8, 10.8 and 10.6 kg CO<sub>2</sub> eq. ha<sup>-1</sup> yr<sup>-1</sup> in GL0-CC, GL1-CC and GL2-CC, respectively. The contribution of indirect emissions of N<sub>2</sub>O from nitrate leaching and ammonia volatilization on the C footprint excluding SOC was low, representing less than a 2% of the total emissions.

### 3.3.3. *Changes in soil organic carbon stocks.*

The annual change in the stock of soil organic carbon (SOC) at the plow layer (0-30 cm depth) was reported in a previous publication (Plaza-Bonilla *et al.*, 2016). With the exception of the GL0-CC cropping system, which sequestered SOC at a rate of 284 kg C ha<sup>-1</sup> yr<sup>-1</sup>, the rest of cropping systems significantly lost SOC. The loss was of a greater magnitude when grain legumes were included in the cropping system, although mitigated with the introduction of cover crops: -233, -595 and -735 kg C ha<sup>-1</sup> yr<sup>-1</sup> in GL0-BF, GL1-BF and GL2-BF, respectively, and -493 and -246 in GL1-CC and GL2-CC, respectively.

### 3.4. *Cropping systems carbon footprint and its relation to grain C.*

The change in SOC stocks had a great and key impact on the C footprint of the different cropping systems (Table 6). Carbon footprint of the six cropping systems showed a high variation with values ranging between 977 and 4015 kg CO<sub>2</sub>eq ha<sup>-1</sup> yr<sup>-1</sup> (Table 6), being found these values in the GL0-CC and the GL2-BF treatments, respectively, when including SOC in the assessment. When no cover crops during the fallow period were cropped between main cash crops, the increase in the number of grain legumes in the 3-yr rotations led to a significant increase in the C footprint. Thanks to a longer photosynthetic activity at the rotation scale and then a greater CO<sub>2</sub> capture, the introduction of cover crops led to a significant decrease in the carbon footprint in the rotations without and with two grain legumes, compared with their counterparts without cover crops (Table 6). When SOC stocks were excluded from the quantification of C footprint the cropping systems without grain legumes (GL0-BF and GL0-CC) showed significantly greater values than the ones with grain legumes (Table 6). Moreover, in this case, the introduction of cover crops increased the C footprint in the cropping systems with one and two grain legumes (i.e. GL1-CC > GL1-BF and GL2-CC > GL2-BF). The differences found between cropping systems on the quotient between C footprint including SOC and the amount of grain C produced were in the line of the ones obtained in the C footprint (Table 6). Due to lower yields of grain legume than cereals, the cropping systems with one grain legume (GL1-BF and GL1-CC) and the one with two grain legumes without cover crops (GL2-BF) presented the highest C footprint per each kg of grain C produced, with values between 2100 and 2430 g CO<sub>2</sub> eq kg<sup>-1</sup> grain C. When grain N was used in the quotient, GL2-BF showed lower values than GL1-BF and GL1-CC, thanks to its greater protein production. Finally, SOC exclusion lowered the ratios between C footprint and grain C and N in the cropping systems with grain legumes. Therefore, the introduction of cover crops lowered the C footprint per each kg of grain C and per each kg of grain N produced in the rotations without and with two grain legumes.

## 4. Discussion

### 4.1. SOC changes as a key component for analyzing multi-year C footprint.

The objective of this study was to analyze the impact of the diversification of cropping systems with grain legumes and cover crops on the C footprint up to the farm gate considering two entire 3-year rotation cycles. This duration allowed obtaining significant conclusions corresponding to the pedoclimatic zone, which presents inter-annual weather variability. Given the cropping system framework of analysis (Drinkwater, 2002), not only the crop sequence differed between treatments but also different crop management practices were adapted. The quantification of the C footprint up to the farm gate for different cropping systems encompasses important items of the ‘farming systems’ framework. Among other, (i) our study compares different orientation of arable crop farms and (ii) uses C footprint as a holistic indicator, which integrates fieldwork (i.e. machinery use), crop productivity and several environmental impacts of cropping practices.

One of the main outcomes of this study is the great impact of SOC changes on the magnitude of C footprint of the different cropping systems analyzed. When SOC change was excluded from the quantification, C footprint was lower in the cropping systems including grain legumes compared to GL0. Similarly, Barton *et al.* (2014) observed a 56% reduction in GHG emissions from wheat production when incorporating a lupin (*Lupinus angustifolius* L.) crop in a 2-yr rotation. However, our results indicated different trends when including the change in SOC in the analysis. Some studies pointed out a reduced SOC loss in legume-based cropping systems compared to cereal-based cropping systems (Drinkwater *et al.*, 1998), mainly when legume leys are established, since they allocate more carbon below-ground and drastically reduce bare fallow periods (Freibauer *et al.*, 2004). In the present study, greater losses of SOC were observed when introducing annual grain legumes in the 3-yr rotations without cover crops. Differently, the rotation without grain legumes and with cover crops (GL0-CC) showed an

increase in SOC stocks after 6 years which reduced significantly its C footprint. This result would be the consequence of the combination of (i) a greater amount of inputs of (ii) lower decomposability and (iii) the possible role played by N limitation in GL1 and GL2. In this line, as an average of the 6 years of experiment, carbon inputs were 2385 and 2978 kg C ha<sup>-1</sup> yr<sup>-1</sup> in the cropping systems without cover crops and with cover crops, respectively (Plaza-Bonilla *et al.*, 2016). Increasing the amount of residue returned to the soil leads to greater SOC stocks (Huggins *et al.*, 1998). In turn, C:N ratio of the crop residues was higher in the cropping systems without grain legumes (62 and 59 for GL0-BF and GL0-CC, respectively, compared to 47, 39, 50 and 41 for GL1-BF, GL1-CC, GL2-BF and GL2-CC, respectively) (Plaza-Bonilla *et al.*, 2016). Moreover, grain legume crop residues have a lower lignin content, which can increase decomposition (Tian *et al.*, 1992). The level of N fertilizer applied in each cropping system could also explain the greater SOC in GL0. Regarding to this, the “microbial nitrogen mining” hypothesis indicates a decrease in soil organic matter decomposition rates at higher nitrogen availability. Thus, in systems with lower availability of N, microorganisms would use labile C to decompose recalcitrant organic matter to acquire N (Craine *et al.*, 2007). Moreover, it has been shown that the application of high levels of N reduces the mass loss of high-lignin materials (Fog, 1988; Knorr *et al.*, 2005) such as sorghum residues, slowing down the overall decomposition process (Mary *et al.*, 1996). In this line, Fog (1988) concluded that N negative effect is mainly found with recalcitrant organic matter of a high C:N ratio, while a positive effect (i.e. greater decomposition with N availability) is commonly found for easily degradable materials. Therefore, the higher N fertilizer rates applied in the GL0 cropping systems in combination with the presence of lignin compounds in sorghum could also partly explain the lower loss of SOC in GL0-BF and SOC increase in GL0-CC compared to the rest of cropping systems.

Depending on their management, agricultural soils can act as a source or sink of CO<sub>2</sub> (i.e. soil carbon sequestration), which is also well illustrated by our results. The impact of agricultural practices on SOC will then increase or offset the CO<sub>2</sub> eq emitted because of crop production activities. In this line, our results highlight the need to take into account SOC changes in carbon footprint analysis in agricultural systems. This aspect agrees with the findings of Yang *et al.* (2014) who observed a significant increase in SOC sequestration (0-20 cm depth) in five diversified crop rotations in the North China Plain, which modified the C footprint of the cropping systems. Similarly, Gan *et al.* (2012) highlighted the significant influence of long-term (25 years) soil carbon change on the value of C footprint for spring wheat when comparing different fallow frequencies in Saskatchewan, Canada. The last authors reported a reversion of the carbon footprint values from positive to negative (i.e. soils acting as a net sink of CO<sub>2</sub> counteracting the CO<sub>2</sub> eq emitted by the rest of components within the C footprint thresholds) when including SOC changes in the calculation, as also stressed by Knudsen *et al.* (2014). However, it must be taken into account that SOC levels could reach an apparent equilibrium after any change in the cropping system (management practices and/or crop rotation) (West and Six, 2007). The timeframe needed to reach this equilibrium will depend on soil management, historical land-use and pedoclimatic conditions (West *et al.*, 2004). Then, although our data indicates a great impact of grain legumes and cover crops on SOC in the short-term, this process would have a lower impact when reaching the new steady state. Differently, other items of the C footprint such as direct and indirect GHG emissions from N fertilizers would be permanently affected by the adoption of this novel cropping systems. Therefore, fostering grain legumes would reduce the C footprint in the long-term.

#### *4.2. Grain legumes and cover crop incorporation in the cropping systems: impacts on C footprint.*

The introduction of grain legumes in the crop rotations without cover crops (GL1-BF and GL2-BF) led to a greater C footprint, which was not necessarily expected. Nevertheless, Jensen *et al.* (2012) suggested a potential SOC sequestration when including grain legumes in cropping systems, provided reduced tillage or no-tillage techniques are used. Therefore, one possible strategy to reduce the negative impact of SOC losses on C footprint in the cropping systems with legumes tested could be the adoption of reduced tillage or no-tillage practices that should be compatible to low input systems in order to avoid a greater use of pesticides.

The introduction of grain legumes led to a significant decrease in the external emissions related to fertilizers, mainly due to the lower requirements of N (as ammonium nitrate and urea) associated to the biological nitrogen fixation of the grain legumes. The manufacturing of nitrogen fertilizer is a highly energy-consuming process, which relies on high temperatures and pressures to synthesize  $\text{NH}_3$ , resulting in a significant amount of  $\text{CO}_2$  emitted to the atmosphere (Nemecek and Erzinger, 2005). Moreover, the use of a lower amount of urea fertilizer to sunflower in GL1 also led to a reduction in the amount of  $\text{CO}_2$  volatilized from urea hydrolysis. The complete reliance of the cropping system without legumes (GL0-BF) on N fertilizer also led to greater on-site emissions as direct soil  $\text{N}_2\text{O}$  emissions, as simulated with the STICS model. According to the model, these emissions would be greatest during N fertilizer applications and, in specific cases, when incorporating low C:N ratio biomass from legume cover crops, which agrees with the existing literature (Bouwman *et al.*, 2013; Rochette and Janzen, 2005). The magnitudes simulated with the model are in agreement with the values reported by Peyrard *et al.* (2016) and Plaza-Bonilla *et al.* (2017a), who measured and simulated the emission of  $\text{N}_2\text{O}$  on a novel version of this cropping system experiment also including legumes, such as faba bean and cover crop mixtures. Nitrous oxide is a powerful greenhouse gas, resulting from a range of biological N transformation processes, being the nitrification and denitrification of a great importance (Butterbach-Bahl *et al.*, 2013). The introduction of

legumes in crop rotations has been long discussed as a key strategy to reduce the direct emissions of N<sub>2</sub>O from soils, due to the reduction of N fertilizer needs (Jensen *et al.*, 2012). Improving N use efficiency in arable cropping systems is key to improve their environmental performance (Prechsl *et al.*, 2017).

The introduction of cover crops in the cropping systems was crucial to reduce the C footprint. The use of cover crops in cropping systems is a promising strategy to sequester carbon in soils, among other benefits, since the period of photosynthesis is lengthened at the rotation scale. In this line, in a meta-analysis, Poeplau and Don (2015) reported an annual SOC sequestration rate of 0.32 Mg ha<sup>-1</sup> when using cover crops. However, according to the results, the introduction of cover crops slightly increased the external emissions related to fertilizers. In our experiment, the N fertilizer rate was slightly increased in the cropping systems with cover crops according to the results of the balance-sheet methodology used to define N fertilization (Plaza-Bonilla *et al.*, 2017b). Cover crop management was optimized with an early date of incorporation in November (Table 5). This early date was determined to avoid preemptive competition for water and N and have a good synchrony of N release from the decomposition of cover crops residues with the needs of the subsequent cash crop. Preemptive competition stands for the reduction of the mineral N caused by cover crop use, reducing the amount of N available for the succeeding crop. This process mainly occurs in seasons with low or no water drainage and concomitant low or no nitrate leaching (Thorup-Kristensen and Nielsen, 1998), as occurred in the experimental area in the first four cropping seasons, therefore, N fertilization needs to be adapted.

The cropping systems with cover crops also presented greater on-site emissions related to the machinery use for their establishment and termination (i.e. pre-sowing soil tillage, sowing and cover crops incorporation to the soil). Similarly, Prechsl *et al.*, (2017) reported slightly higher global warming potential due to the energy demand in cropping systems with cover



506 crops compared to the use of fallow between cash crops in a cropping systems experiment in  
507 Switzerland. These emissions did not counteract the lower C footprint in these cropping  
508 systems, which was the result of the positive incidence of cover crops on the mitigation of SOC  
509 losses. However, the energy needs and concomitant costs resulting from cover crops use can  
510 suppose a hindrance for the implementation of this strategy by farmers (Gabriel *et al.*, 2013).

511

## 5. Conclusions

The results of this study highlighted the importance of including soil organic carbon changes for doing a relevant analysis in C footprint assessments of arable cropping systems. Grain legumes introduction into low input crop rotations led to SOC losses, counteracting the positive impact of legumes in the reduction in external and on-site emissions related to N fertilizers. Therefore, our results support the need to include SOC changes in environmental assessments of agricultural production. However, according to the literature, SOC changes are finite in time if the different factors regulating the soil C balance (e.g. crop management practices, crop sequence, climatic conditions) remain unchanged. Therefore, the timeframe of environmental assessments of cropping systems must be clearly identified. Cover crops introduction in the cropping systems analyzed was key to reduce the C footprint, due their positive impact on SOC loss mitigation, indicating their potential as a CO<sub>2</sub> offsetting strategy in agricultural production. However, cover crops need to be carefully implemented by farmers given the economic costs associated to management practices (i.e. establishment and termination). It can then be recommended to policy makers that the use of cover crops must be explicitly taken into account and thus remunerated at their fair value. More research is needed to design grain legume-based cropping systems which allow maintaining SOC levels in low-input agricultural systems to improve the C footprint.

## Acknowledgements

The constructive suggestions of three anonymous reviewers greatly improved this manuscript. This research was supported by (i) the FP6 Grain Legumes Integrated Project (food-CT-2004-506223), (ii) the European Commission (REA) through the LEGATO project (FP7-613551) and (iii) the French National Research Agency (ANR) through the LEGITIMES French project

536 (ANR-13-AGRO-0004). It also benefited from the support of the Climate-CAFE European  
537 project, which was selected by the European FACCE-JPI ERA-NET Plus program. The authors  
538 thank Jean-Marie Nolot for his key role in managing the cropping system experiment of  
539 Auzeville and Grégory Véricel for the database creation and maintenance. INS was funded  
540 thanks to a convention between the University of Lleida and INRA-Toulouse. DPB received a  
541 Juan de la Cierva postdoctoral grant from the Ministerio de Economía y Competitividad of  
542 Spain (IJCI-2016-27784).

543

## References

- ADEME (Agence de l'Environnement et de la Maîtrise de l'Energie). 2011. Guide des valeurs  
Dia'terre. Version référentiel 1.13. (In French). 187 p. Angers, France.
- Audsley, E.; Stacey, K.; Parsons, D.J.; Williams, A.G. 2009. Estimation of the greenhouse gas  
emissions from agricultural pesticide manufacture and use. Cranfield University.
- Barton, L.; Thamo, T.; Engelbrecht, D.; Biswas, W.K. 2014. Does growing grain legumes or  
applying lime cost effectively lower greenhouse gas emissions from wheat production in  
semi-arid climate? *Journal of Cleaner Production* 83, 194-203.
- Bedoussac, L.; Journet, E-P.; Hauggaard-Nielsen, H.; Naudin, C.; Corre-Hellou, G.; Jensen,  
E.S.; Prieur, L.; Justes, E. 2015. Ecological principles underlying the increase of  
productivity achieved by cereal-grain legume intercrops in organic farming. A review.  
*Agronomy for Sustainable Development* 35, 911-935.
- Bouwman, A.F.; Beusen, A.H.W.; Griffioen, J.; Van Groenigen, J.W.; Hefting, M.M.; Oenema,  
O.; Van Puijenbroek, P.J.T.M.; Seitzinger, S.; Slomp, C.P.; Stehfest, E. 2013. Global trends  
and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philosophical  
Transactions of the Royal Society B Biological Sciences* 368, 20130112.
- Brisson, N., Mary, B., Ripoche, D., Jeuffroy, M.H., Ruget, F., Nicoullaud, B., Gate, P.,  
Devienne-Barret, F., Antonioletti, R., Durr, C., Richard, G., Beaudoin, N., Recous, S.,  
Tayot, X., Plenet, D., Cellier, P., Machet, J.M., Meynard, J.M., Delecolle, R., 1998. STICS:  
a generic model for the simulation of crops and their water and nitrogen balances. I. Theory  
and parameterization applied to wheat and corn. *Agronomie* 18, 311-346.
- Brisson, N., Ruget, F., Gate, P., Lorgeau, J., Nicoullaud, B., Tayot, X., Plenet, D., Jeuffroy,  
M.H., Bouthier, A., Ripoche, D., Mary, B., Justes, E., 2002. STICS: a generic model for

567       simulating crops and their water and nitrogen balances. II. Model validation for wheat and  
568       maize. *Agronomie* 22, 69-92.

569   Brisson, N., Launay, M., Mary, B., Beaudoin, N. 2008. Conceptual basis, formalisations and  
570       parameterization of the STICS crop model. Editions QUAE, INRA, 78026 Versailles  
571       Cedex.

572   Butterbach-Bahl, K.; Baggs, E.; Dannenmann, M.; Kiese, R.; Zechmeister-Boltenstern, S.  
573       2013. Nitrous oxide emissions from soils: how well do we understand the processes and  
574       their controls? *Philosophical Transactions of the Royal Society B Biological Sciences* 368,  
575       20130122.

576   Ceschia, E.; Béziat, P.; Dejoux, J-F.; Aubinet, M.; Bernhofer, C. *et al.* 2010. Management  
577       effects on net ecosystem carbon and GHG budgets at European crop sites. *Agriculture,*  
578       *Ecosystems and Environment* 139, 363-383.

579   Craine, J.M.; Morrow, C.; Fierer, N. 2007. Microbial nitrogen limitation increases  
580       decomposition. *Ecology* 88, 2105-2113.

581   Drinkwater, L.E. 2002. Cropping systems research: reconsidering agricultural experimental  
582       approaches. *HortTechnology* 12, 355-361.

583   Drinkwater, L.E.; Wagoner, P.; Sarrantonio, M. 1998. Legume-based cropping systems have  
584       reduced carbon and nitrogen losses. *Nature*, 262-265.

585   Fog, K. 1988. The effect of added nitrogen on the rate of decomposition of organic matter.  
586       *Biological Reviews* 63, 433-462.

587   Freibauer, A.; Rounsevell, M.D.A.; Smith, P.; Verhagen, P. 2004. Carbon sequestration in the  
588       agricultural soils of Europe. *Geoderma* 122, 1-23.

589 Gabriel, J.L.; Garrido, A.; Quemada, M. 2013. Cover crops effect on farm benefits and nitrate  
590 leaching: Linking economic and environmental analysis. *Agricultural Systems* 121, 23-32.

591 Gaillard, G.; Hausheer, J.; Crettaz, P. 1997. Inventaire environnemental des intrants agricoles  
592 en production végétale. Base de données pour l'établissement des bilans énergétiques et  
593 écologiques en agriculture. Station fédérale de recherche en économie et technologies  
594 agricoles. Tanikon, Switzerland, 49 pp.

595 Gan, Y.; Liang, C.; Campbell, C.A.; Zentner, R.P.; Lemke, R.L.; Wang, H.; Yang, C. 2012.  
596 Carbon footprint of spring wheat in response to fallow frequency and soil carbon changes  
597 over 25 years on the semiarid Canadian prairie. *European Journal of Agronomy* 43, 175-  
598 184.

599 Huggins, D.R.; Buyanovsky, G.A.; Wagner, G.H.; Brown, J.R.; Darmody, R.G.; Peck, T.R.;  
600 Lesoing, G.W.; Vanotti, M.B.; Bundy, L.G. 1998. Soil organic C in the tallgrass prairie-  
601 derived region of the corn belt: effects of long-term crop management. *Soil and Tillage*  
602 *Research* 47, 219–234.

603 IPCC. 2006. Agriculture, forestry and other land use. In: Eggleston, H.S.; Buendia, L.; Miwa,  
604 K.; Ngara, T.; Tanabe, K. (Eds.), 2006 IPCC guidelines for national greenhouse gas  
605 inventories. Institute for Global Environmental Strategies, Hayama, Japan.

606 IPCC. 2013. Contribution of working group I to the fifth assessment report of the  
607 Intergovernmental Panel on Climate Change. *Climate change 2013: The physical science*  
608 *basis*. Cambridge University Press, Cambridge, United Kingdom and New York NY, USA.

609 Jambert, C.; Delmas, R.; Serça, D.; Thouron, L.; Labroue, L.; Delprat, L. 1997. N<sub>2</sub>O and CH<sub>4</sub>  
610 emissions from fertilized agricultural soils in southwest France. *Nutrient Cycling in*  
611 *Agroecosystems* 48, 105-114.

612 Jensen, E. S.; Peoples, M.B.; Boddey, R.M.; Gresshoff, P.M.; Hauggaard-Nielsen, H.; Alves,  
613 B.J.R.; Morrison, M.J. 2012. Legumes for mitigation of climate change and the provision  
614 of feedstock for biofuels and biorefineries. A review. *Agronomy for Sustainable*  
615 *Development* 32, 329-364.

616 Knorr, M.; Frey, S.D.; Curtis, P.S. 2005. Nitrogen additions and litter decomposition: a meta-  
617 analysis. *Ecology* 86, 3252-3257.

618 Knudsen, M.T.; Meyer-Aurich, A.; Olesen, J.E.; Chirinda, N.; Hermansen, J.E. 2014. Carbon  
619 footprints of crops from organic and conventional arable crop rotations – using a life cycle  
620 assessment approach. *Journal of Cleaner Production* 64, 609-618.

621 Lal, R. 2004. Carbon emission from farm operations. *Environmental International* 30, 981-990.

622 Lares-Orozco, M.F.; Robles-Morúa, A.; Yepez, E.A.; Handler, R.M. 2016. Global warming  
623 potential of intensive wheat production in the Yaqui Valley, Mexico: a resource for the  
624 design of localized mitigation strategies. *Journal of Cleaner Production* 127, 522-532.

625 Mary, B.; Recous, S.; Darwis, D.; Robin, D. 1996. Interactions between decomposition of plant  
626 residues and nitrogen cycling in soil. *Plant and Soil* 181, 71-82.

627 Nemecek, T.; Erzinger, S. 2005. Modelling representative life cycle inventories for Swiss  
628 arable crops. *The International Journal of Life Cycle Assessment* 10, 68-76.

629 Pandey, D.; Agrawal, M. 2014. Carbon footprint estimation in the agriculture sector. In: Muthu,  
630 S.S. (Ed.) *Assessment of carbon footprint in different industrial sectors. Volume I.*  
631 *EcoProduction*. Springer, Singapore. ISBN 978-981-4560-40-5.

632 Pandey, D.; Agrawal, M.; Pandey, J.S. 2011. Carbon footprint: current methods of estimation.  
633 *Environmental Monitoring and Assessment* 178, 135-160.

634 Peyrard, C.; Mary, B.; Véricel, G.; Gréhan, E.; Justes, E.; Léonard, J. 2016. N<sub>2</sub>O emissions of  
635 low input cropping systems as affected by legume and cover crops use. *Agriculture,*  
636 *Ecosystems and Environment* 224, 145-156.

637 Plaza-Bonilla, D.; Léonard, J.; Peyrard, C.; Mary, B.; Justes, E. 2017a. Precipitation gradient  
638 and crop management affect N<sub>2</sub>O emissions: Simulation of mitigation strategies in rainfed  
639 Mediterranean conditions. *Agriculture, Ecosystems and Environment* 238, 89-103.

640 Plaza-Bonilla, D.; Nolot, J-M.; Passot, S.; Raffaillac, D.; Justes, E. 2016. Grain legume-based  
641 rotations managed under conventional tillage need cover crops to mitigate soil organic  
642 matter losses. *Soil and Tillage Research* 156, 33-43.

643 Plaza-Bonilla, D.; Nolot, J-M.; Raffaillac, D.; Justes, E. 2015. Cover crops mitigate nitrate  
644 leaching in cropping systems including grain legumes: Field evidence and model  
645 simulations. *Agriculture, Ecosystems and Environment* 212, 1-12.

646 Plaza-Bonilla, D.; Nolot, J-M.; Raffaillac, D.; Justes, E. 2017b. Innovative cropping systems  
647 to reduce N inputs and maintain wheat yields by inserting grain legumes and cover crops  
648 in southwestern France. *European Journal of Agronomy* 82, 331-341.

649 Poeplau, C.; Don, A. 2015. Carbon sequestration in agricultural soils via cultivation of cover  
650 crops – A meta-analysis. *Agriculture, Ecosystems and Environment* 200, 33-41.

651 Prechsl, U.E.; Wittwer, R.; van der Heijden, M.G.A.; Lüscher, G.; Jeanneret, P.; Nemecek, T.  
652 2017. Assessing the environmental impacts of cropping systems and cover crops: Life cycle  
653 assessment of FAST, a long-term arable farming field experiment. *Agricultural Systems*  
654 157, 39-50.



655 Reckling, M.; Bergkvist, G.; Watson, C.A.; Stoddart, F.L.; Zander, P.M.; Walker, R.L.; Pristeri,  
656 A.; Toncea, I.; Bachinger, J. 2015. Trade-offs between economic and environmental  
657 impacts of introducing legumes into cropping systems. *Frontiers in Plant Science* 7:669.

658 Rochette, P.; Janzen, H.H. 2005. Towards a revised coefficient for estimating N<sub>2</sub>O emissions  
659 from legumes. *Nutrient Cycling in Agroecosystems* 73, 171-179.

660 SAS Institute Inc, 2017. Using JMP®13 Cary, NC: SAS Institute Inc.

661 Thorup-Kristensen, K.; Nielsen, N.E. 1998. Modelling and measuring the effect of nitrogen  
662 catch crops on the nitrogen supply for succeeding crops. *Plant and Soil* 203, 79-89.

663 Tian, G.; Kang, B.T.; Brussaard, L. 1992. Biological effects of plant residues with contrasting  
664 chemical compositions under humid tropical conditions-decomposition and nutrient  
665 release. *Soil Biology and Biochemistry* 24, 1051-1060.

666 Watson, C.A.; Reckling, M.; Preissel, S.; Bachinger, J.; Bergkvist, G.; Kuhlman, T.; Lindström,  
667 K.; Nemecek, T.; Topp, C.F.E.; Vanhatalo, A.; Zander, P.; Murphy-Bokern, D.; Stoddart,  
668 F.L. 2017. Grain legume production and use in European agricultural systems. *Advances*  
669 *in Agronomy* 144, 235-303.

670 West, T.O.; Marland, G.; King, A.W.; Post, W.M.; Jain, A.K.; Andrasko, K. 2004. Carbon  
671 management response curves: estimates of temporal soil carbon dynamics. *Environmental*  
672 *Management* 33, 507-518.

673 West, T.O.; Six, J. 2007. Considering the influence of sequestration duration and carbon  
674 saturation on estimates of soil carbon capacity. *Climatic Change* 80, 25-41.

675 Yang, X.; Gao, W.; Zhang, M.; Chen, Y.; Sui, P. 2014. Reducing agricultural carbon footprint  
676 though diversified crop rotation systems in the North China Plain. *Journal of Cleaner*  
677 *Production* 76, 131-139.

## Figure captions

**Figure 1** Cropping systems studied in the field experiment. GL0, GL1 and GL2 stand for three-year rotation with 0, 1, and 2 grain legumes. Cash crops (big circles) and cover crops (small circles) are shown. For color version of this figure, the reader is referred to the online version of this article.

**Figure 2** Monthly precipitation (grey bars) and air temperature (solid black line) at the experimental site: 30-yr (1981-2010) average values and 2004-2005, 2005-2006, 2006-2007, 2007-2008, 2008-2009 and 2009-2010 cropping seasons. Cropping season precipitation is shown in italics.

**Figure 3** External emissions related to fertilizer, pesticides and seeds production and transportation and to energy needed to pump irrigation water as affected by cropping system (GL0, GL1 and GL2, 0, 1 and 2 grain legumes in a 3-years rotation; CC, cover crop and BF, bare fallow). Values correspond to the annual average of the 2005 – 2010 period. Vertical bars correspond to the standard deviation. For color version of this figure, the reader is referred to the online version of this article.

**Figure 4** On-site emissions related to machinery use, N<sub>2</sub>O indirect emissions as nitrate leaching and ammonia volatilization, soil N<sub>2</sub>O direct emissions, and CO<sub>2</sub> emissions due to urea hydrolysis as affected by cropping system (GL0, GL1 and GL2, 0, 1 and 2 grain legumes in a 3-years rotation; CC, cover crop and BF, bare fallow). Values correspond to the annual average of the 2005 – 2010 period. Vertical bars correspond to the standard deviation. For color version of this figure, the reader is referred to the online version of this article.

**Figure 5** Simulated soil N<sub>2</sub>O emissions (2004–2010) as affected by crop rotation (GL0, GL1 and GL2, 3-yr rotations with 0, 1 and 2 grain legumes, respectively) and cover crop treatments. The three crop sequences of each rotation are shown horizontally as sub-figures (a, b and c:

702 crop sequences of GL0; d, e and f, crop sequences of GL1; g, h and I, crop sequences of GL2).  
703 Dashed lines delimitate the season covered by each cash (in black letters) and cover crop (in  
704 grey letters) and fallow periods. DW, Mu, SF, sP, SR, Sy, V, VO, wP stand for durum wheat,  
705 mustard, sunflower, spring pea, sorghum, soybean, vetch, vetch-oat mixture and winter pea,  
706 respectively. Note that each sub-figure shares X-axis. For color version of this figure, the reader  
707 is referred to the online version of this article.

708

**Table 1.** Average sowing and harvest dates of cash crops and sowing rate ranges for cash and cover crops used in the experiment. In the vetch-oat crop mixture, values refer to the crop between brackets.

Crop	Sowing date	Harvest date	Seeding rate (seeds m <sup>-2</sup> )
Durum wheat	November	July (first fortnight)	278-413
Winter pea	December	June (last fortnight)	70-93
Spring pea	February	June (last fortnight)	100-113
Sorghum	end April – beginning May	September (last fortnight)	29-32
Sunflower	April (last fortnight)	September	6.6-6.7
Soybean	May (first week)	end September – beginning October	36-39
(Vetch)-Oat	After cash crop harvest	Not harvested	31-83
Vetch-(Oat)	After cash crop harvest	Not harvested	43-170
Vetch	After cash crop harvest	Not harvested	50-76
Mustard	After cash crop harvest	Not harvested	135-172

713 **Table 2.** Nitrogen fertilizer rates applied to the different cash crops in the cropping systems studied (GL0, GL1 and GL2, 3-year rotation with 0,  
 714 1 and 2 grain legumes, respectively; BF and CC, bare fallow and cover crops, respectively). Splits of N to durum wheat are shown. Note that no  
 715 N fertilizer was applied to grain legumes and cover crops.

Rotation	Cash crop	N fertilizer application (kg N ha <sup>-1</sup> )													
		Bare fallow (BF)							Cover crop (CC)						
		2005	2006	2007	2008	2009	2010	Mean	2005	2006	2007	2008	2009	2010	Mean
GL0	Durum wheat	132+50	99+50	100+54	101+57	42+74+55	100+80	166	88+50	99+50	100+54	101+57	42+74+55	100+80	158
	Sorghum	76	76	82	83	112	60	82	76	76	82	83	112	60	82
	Sunflower	56	51	51	62	67	40	55	56	51	51	62	67	40	55
	3-yr N applied							303							295
GL1	Durum wheat	66+50	50+50	50+54	40+57	106+55	60+80	120	66+50	99+50	100+54	70+57	106+55	80+80	145
	Sunflower	0	0	0	0	34	0	6	0	0	0	0	34	0	6
	Winter pea	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	3-yr N applied							128							151
GL2	Durum wheat	88+50	50+50	50+54	40+57	42+37+55	80+50	117	88+50	99+50	50+54	70+57	42+56+55	60+50	130
	Soybean	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Spring pea	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	3-yr N applied							117							130

716

717 **Table 3** Emission factors (EF) for the different inputs used in the cropping systems experiment.

Input type	Description	EF	Source
Fertilizer	Ammonium nitrate	1.11 kg Ceq kg <sup>-1</sup>	Ceschia <i>et al.</i> (2010)
	Urea	1.29 kg Ceq kg <sup>-1</sup>	
	Triple superphosphate	0.42 kg Ceq kg <sup>-1</sup>	
Herbicides	Different active ingredients	Range: 0.41-5.80 kg Ceq kg <sup>-1</sup> a.i.	Audsley <i>et al.</i> (2009) Lal (2004)
Fungicides	Different active ingredients	Range: 0.55-1.95 kg Ceq kg <sup>-1</sup> a.i.	Audsley <i>et al.</i> (2009) Gaillard <i>et al.</i> (1997)
Insecticides	Different active ingredients	Range: 1.54-1.90 kg Ceq kg <sup>-1</sup> a.i.	Audsley <i>et al.</i> (2009)
Irrigation	Energy for pumping water	0.516 kg Ceq mm <sup>-1</sup>	Ceschia <i>et al.</i> (2010)
Seeds		0.04-0.21 kg Ceq kg <sup>-1</sup>	ADEME (2011)

718

**Table 4** Farming operations carried out in the field experiment, and associated tools/implements, and fuel consumption.

Farming operation	Tool/Implement	Fuel consumption (L ha <sup>-1</sup> )
Fertilization	Pneumatic spreader 800 L	1.5
Harvest	Medium-sized combine	18
Irrigation	Hose reel irrigation system	-
Mechanical weed control	Cultivator (camera steering system)	5
Pesticides application	Sprayer	2
Residues chopping	Shredder	10
Rolling	Cultipacker	3
Sowing – Small grains	Seed driller	7
Sowing – Coarse grains	Pneumatic row planter	4.5
Tillage – Cultivation	Cultivator	12
Tillage – Harrowing	Disk harrow – 22 disks	10
Tillage – Germinator	Tine harrow	6.5
Tillage – Plowing	Moldboard plow	27
Tillage – Harrowing	Rotary Harrow	13
Tillage – Subsoiling	Subsoiler	17
Tillage – Shallow cultivation	Vibro-cultivator	6



**Table 5** Cash crops grain yield (dry matter) in the different cropping systems compared (GL0, GL1 and GL2, 3-year rotation with 0, 1 and 2 grain legumes, respectively; BF and CC, bare fallow and cover crops, respectively). Values correspond to the average of 6 years (i.e. 2005-2010 period). Values between brackets correspond to the standard deviation.

Rotation	Cash crop	Cover crop treatment	
		BF	CC
GL0	Sorghum	6613 (617)	6782 (709)
	Sunflower	2678 (455)	2506 (463)
	Durum wheat	4974 (799)	5012 (803)
GL1	Sunflower	2581 (599)	2693 (473)
	Winter pea	2645 (1316)	2705 (1027)
	Durum wheat	5458 (902)	5296 (666)
GL2	Soybean	3092 (635)	3005 (726)
	Spring pea	3260 (1123)	3322 (1127)
	Durum wheat	5282 (734)	4991 (1120)


729 **Table 6** Carbon footprint, C footprint per kg of grain carbon, and C footprint per kg of grain nitrogen including and excluding soil organic carbon  
730 (SOC) changes as affected by cropping system (GL0, GL1 and GL2, 3-year rotation with 0, 1 and 2 grain legumes, respectively; BF and CC, bare  
731 fallow and cover crops, respectively). Values correspond to the annual average of the 2005 – 2010 period. Values between brackets correspond to  
732 the standard deviation. Different letters indicate differences between cropping systems at  $P<0.05$ .

Cropping system	C footprint (kg CO <sub>2</sub> eq ha <sup>-1</sup> yr <sup>-1</sup> )		C footprint kg <sup>-1</sup> grain C (g CO <sub>2</sub> eq kg <sup>-1</sup> grain C)		C footprint kg <sup>-1</sup> grain N (kg CO <sub>2</sub> eq kg <sup>-1</sup> grain N)	
	Including SOC	Excluding SOC	Including SOC	Excluding SOC	Including SOC	Excluding SOC
GL0-BF	2897 (111) c	2043 (111) a	1423 (89) b	974 (67) a	32 (4) b	22 (3) a
GL0-CC	977 (101) e	2018 (102) a	411 (57) c	976 (77) a	9 (1) d	21 (2) a
GL1-BF	3344 (127) b	1162 (127) d	2430 (709) a	698 (140) b	41 (12) a	12 (3) bc
GL1-CC	3282 (142)b	1475 (142) b	2100 (252) a	824 (48) b	36 (5) ab	14 (1) b
GL2-BF	4015 (44) a	1320 (44) c	2458 (413) a	744 (103) b	31 (6) b	10 (2) c
GL2-CC	2389 (139) d	1487 (139) b	1429 (182) b	842 (122) ab	18 (3) c	11 (2) bc

733




**Figure 1**




737

738 **Figure 2**




**Figure 3**





**Figure 4**



**Figure 5**

